



**Europäisches  
Patentamt**

**European  
Patent Office**

**Office européen  
des brevets**

REC'D 24 AUG 2004

WIPO

PCT

**RECEIVED :**

05 JUL. 2004

Patent & Licensing Dept.  
BAYER CROPSCIENCE S.A.

**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

**Patentanmeldung Nr. Patent application No. Demande de brevet n°**

03356065.7

BCS 03.4004

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**R C van Dijk**



Anmeldung Nr:  
Application no.: 03356065.7  
Demande no:

Anmeldetag:  
Date of filing: 15.04.03  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Bayer CropScience SA  
55, avenue René Cassin,  
CP 106  
69266 Lyon Cedex 09  
FRANCE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Fungicidal composition comprising a pyridylmethylbenzamide derivative and a  
chloronitrile derivative

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

A01N/

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT RO SE SI SK TR LI

**Fungicidal composition comprising a pyridylmethylbenzamide derivative and a chloronitrile derivative**

5           The present invention relates to novel fungicidal compositions comprising at least a pyridylmethylbenzamide derivative and a chloronitrile derivative. The present invention also relates to a method of combating phytopathogenic fungi by applying at a locus infested or liable to be infested such a composition.

10           European patent application EP-A-1056723 generically discloses the possibility of combining pyridylmethylbenzamide derivatives with known fungicidal products to develop a fungicidal activity, without citing by name examples of co-active ingredients, or even families of co-active ingredients, which are capable of being combined with pyridylmethylbenzamide derivatives.


15           International patent application WO 02/069713 discloses fungicidal mixtures comprising a pyridylmethylbenzamide derivative and phosphorous acid or one of its derivatives. No mention is made in this patent application of a mixture comprising a pyridylmethylbenzamide derivative and a chloronitrile derivative.

20           Some of the above mentioned mixtures have shown a synergistic effect. Nevertheless, it is always of high-interest in agriculture to use novel pesticidal mixtures showing a synergistic effect in order to avoid or to control the development of resistant strains to the active ingredients or to the mixtures of known active ingredients used by the farmer while minimising the doses of chemical products spread in the environment and reducing the cost of the treatment.

25           We have now found some novel fungicidal compositions which possess the above mentioned characteristics.

          Accordingly, the present invention relates to a fungicidal composition comprising :

30       a) at least a pyridylmethylbenzamide derivative of general formula (I)



in which :

- R<sup>1</sup> may be a hydrogen atom, an optionally substituted alkyl group or an optionally substituted acyl group;

- R<sup>2</sup> may be a hydrogen atom or an optionally substituted alkyl group;

5        - R<sup>3</sup> and R<sup>4</sup> may be chosen independently from each other as being a halogen atom, a hydroxyl group, a cyano group, a nitro group, -SF<sub>5</sub>, a trialkylsilyl group, an optionally substituted amino group, an acyl group, or a group E, OE or SE, in which E may be an alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl or a heterocyclyl group each of which may optionally be substituted;

10        - p represents 0, 1, 2, 3 or 4;

- q represents 0, 1, 2, 3 or 4;

and its agriculturally acceptable optical and/or geometric isomers, tautomers and addition salts with an acid or a base;

and

15        b) a chloronitrile derivative which is chlorothalonil;

in a compound (I) / chlorothalonil weight ratio of from 0,01 to 1.

Chlorothalonil is a fungicide compound also known under its chemical name which is tetrachloroisophthalonitrile.

20        In the context of the present invention :

- the term halogen means bromine, chlorine, iodine or fluorine;

- the term alkyl means a linear or branched saturated hydrocarbon group containing from 1 to 6 carbon atoms;

25        - the term alkenyl means a linear or branched hydrocarbon group containing from 2 to 6 carbon atoms and an unsaturation in the form of double bond;

- the term alkynyl means a linear or branched hydrocarbon group containing from 2 to 6 carbon atoms and an unsaturation in the form of a triple bond;

- the term alkoxy means a linear or branched alkyloxy group containing from 1 to 6 carbon atoms;

30        - the term acyl means a formyl group or linear or branched alkoxycarbonyl group containing from 2 to 6 carbon atoms;

- the term cycloalkyl means a saturated cyclic hydrocarbon group containing from 3 to 8 carbon atoms;

- the term aryl means a phenyl or naphthyl group;

35        - the term heterocyclyl means saturated, partially saturated, unsaturated or aromatic cyclic group containing from 3 to 8 atoms, which may be a carbon atom, a nitrogen

atom, a sulphur atom or an oxygen atom. Examples of such heterocyclyl may be pyridyl, pyridinyl, quinolyl, furyl, thienyl, pyrrolyl, oxazolinyl;

- the term "optionally substituted " means that the group thus termed may be substituted with one or more groups which may be halogen, alkyl, alkoxy, hydroxyl, nitro, amin, cyano or acyl.

The composition according to the present invention provides a synergistic effect. This synergistic effect allows a reduction of the chemical substances spread into the environment and a reduction of the cost of the fungal treatment.

In the context of the present invention, the term "synergistic effect" is defined by Colby according to the article entitled "Calculation of the synergistic and antagonistic responses of herbicide combinations" Weeds, (1967), 15, pages 20-22.

The latter article mentions the formula:

$$E = x + y - \frac{x * y}{100}$$

in which E represents the expected percentage of inhibition of the disease for the combination of the two fungicides at defined doses (for example equal to x and y respectively), x is the percentage of inhibition observed for the disease by the compound (I) at a defined dose (equal to x), y is the percentage of inhibition observed for the disease by the compound (II) at a defined dose (equal to y). When the percentage of inhibition observed for the combination is greater than E, there is a synergistic effect.

The fungicidal mixtures according to the present invention comprises at least a pyridylmethylbenzamide derivative of general formula (I).

Preferably, the present invention relates to a fungicidal mixture comprising at least a pyridylmethylbenzamide derivative of general formula (I) in which the different characteristics may be chosen alone or in combination as being :

- as regards  $R^1$  and  $R^2$ ,  $R^1$  and  $R^2$  are chosen independently from each other as being a hydrogen atom or an optionally substituted alkyl group. More preferably,  $R^1$  and  $R^2$  are chosen independently from each other as being a hydrogen atom, a methyl group or an ethyl group. Even more preferably,  $R^1$  and  $R^2$  are both hydrogen atoms.

- as regards  $R^3$  and  $R^4$ ,  $R^3$  and  $R^4$  are chosen independently from each other as being a halogen atom, a hydroxyl group, a nitro group, an optionally substituted amino group, an acyl group, or a group E, OE or SE, in which E may be a alkyl, a

cycloalkyl, a phenyl or a heterocyclyl group, each of which may optionally be substituted. More preferably,  $R^3$  and  $R^4$  are chosen independently from each other as being a halogen atom, a nitro group or a halogenoalkyl group. Even more preferably  $R^3$  and  $R^4$  are chosen independently from each other as being a chlorine atom, a nitro group or a trifluoromethyl group.

- as regards p, p is 1 or 2. More preferably, p is 2.

- as regards q, q is 1 or 2. More preferably, q is 2;

and its agriculturally acceptable possible tautomers and addition salts with an acid or a base.

More preferably, the present invention relates to a fungicidal mixture comprising at least a pyridylmethylbenzamide derivative of general formula (I) which may be chosen as being :

- a compound (Ia) which is 2,6-dichloro-N-{{3-chloro-5-(trifluoromethyl)-2-pyridinyl}methyl}benzamide; or

- a compound (Ib) which is N-{{3-chloro-5-(trifluoromethyl)-2-pyridinyl}methyl}-2-fluoro-6-nitrobenzamide; or

- a compound (Ic) which is N-{{3-chloro-5-(trifluoromethyl)-2-pyridinyl}methyl}-2-methyl-6-nitrobenzamide;

and its agriculturally acceptable possible tautomers and addition salts with an acid or a base.

The fungicidal mixture according to the present invention comprises at least a pyridylmethylbenzamide derivative of general formula (I) and a chloronitrile derivative which is chlorothalonil in a compound (I) / chlorothalonil weight ratio of from 0,01 to 1; preferably of from 0,025 to 0,5; even more preferably, of from 0,05 to 0,1.

The present invention also relates to a fungicidal composition comprising at least one further active ingredient. Thus, according to the present invention, there is also provided a fungicidal composition as defined above which comprise at least one further fungicidal, herbicidal, insecticide and/or plant growth regulating compounds, according to the use for which they are intended.

The present invention also relates to a fungicidal composition comprising an effective amount of active material and other additional components. Thus, according to the present invention, there is provided a fungicidal composition

comprising active material as defined above and an agriculturally acceptable, support, carrier or filler.

In the present specification, the term "support" denotes a natural or synthetic, organic or inorganic material with which the active material is combined to make it easier to apply, notably to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support may be a solid or a liquid. Examples of suitable supports include clays, natural or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports may also be used.

The composition may also comprise other additional components. In particular, the composition may further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention may be made, for example, of polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulphosuccinic acid esters, taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the above compounds containing sulphate, sulphonate and phosphate functions. The presence of at least one surfactant is generally essential when the active material and/or the inert support are water-insoluble and when the vector agent for the application is water. Preferably, surfactant content may be comprised between 5% and 40% by weight of the composition.

Additional components may also be included, e.g. protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering agents. More generally, the active materials can be combined with any solid or liquid additive, which complies with the usual formulation techniques.

In general, the composition according to the invention may contain from 0.05 to 99% (by weight) of active material, preferably 10 to 70% by weight.

Compositions according to the present invention can be used in various forms such as aerosol dispenser, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, granule, hot fogging concentrate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable

concentrate, oil miscible liquid, paste, plant rodlet, powder for dry seed treatment, seed coated with a pesticide, soluble concentrate, soluble powder, solution for seed treatment, suspension concentrate (flowable concentrate), ultra low volume (ulv) liquid, ultra low volume (ulv) suspension, water dispersible granules or tablets, water  
 5 dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder.

These compositions include not only compositions which are ready to be applied to the plant or seed to be treated by means of a suitable device, such as a spraying or dusting device, but also concentrated commercial compositions which  
 10 must be diluted before they are applied to the crop.

The fungicidal compositions of the present invention can be used to curatively or preventively control the phytopathogenic fungi of crops. Thus, according to a further aspect of the present invention, there is provided a method for preventively or  
 15 curatively controlling phytopathogenic fungi of crops characterised in that a fungicidal composition as hereinbefore defined is applied to the seed, the plant and/or to the fruit of the plant or to the soil in which the plant is growing or in which it is desired to grow.

The composition as used against phytopathogenic fungi of crops comprises an effective and non-phytotoxic amount of an active material of general formula (I).

20 The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention which is sufficient to control or destroy the fungi present or liable to appear on the crops, and which does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the fungus to be combated, the type of crop, the climatic conditions  
 25 and the compounds included in the fungicidal composition according to the invention.

This amount can be determined by systematic field trials, which are within the capabilities of a person skilled in the art.

The method of treatment according to the present invention is useful to treat propagation material such as tubers or rhizomes, but also seeds, seedlings or  
 30 seedlings pricking out and plants or plants pricking out. This method of treatment can also be useful to treat roots. The method of treatment according to the present invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruits of the concerned plant.

Among the plants that can be protected by the method according to the  
 35 invention, mention may be made of cotton; flax; vine; fruit crops such as *Rosaceae* sp. (for instance pip fruits such as apples and pears, but also stone fruits such as



apricots, almonds and peaches), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp., *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp., *Actinidaceae* sp., *Lauraceae* sp., *Musaceae* sp. (for instance banana trees and plantains), *Rubiaceae* sp., *Theaceae* sp., *Sterculiaceae* sp., *Rutaceae* sp. (for instance lemons, oranges and grapefruits); leguminous crops such as *Solanaceae* sp. (for instance tomatoes),  
 5 *Liliaceae* sp., *Asteraceae* sp. (for instance lettuces), *Umbelliferae* sp., *Cruciferae* sp., *Chenopodiaceae* sp., *Cucurbitaceae* sp., *Papilionaceae* sp. (for instance peas), *Rosaceae* sp. (for instance strawberries); big crops such as *Graminae* sp. (for instance maize, cereals such as wheat, rice, barley and triticale), *Asteraceae* sp. (for  
 10 instance sunflower), *Cruciferae* sp. (for instance colza), *Papilionaceae* sp. (for instance soja), *Solanaceae* sp. (for instance potatoes), *Chenopodiaceae* sp. (for instance beetroots); horticultural and forest crops; as well as genetically modified homologues of these crops.

Among the plants and the possible diseases of these plants protected by the  
 15 method according to the present invention, mention may be made of :

- wheat, as regards controlling the following seed diseases: fusaria (*Microdochium nivale* and *Fusarium roseum*), stinking smut (*Tilletia caries*, *Tilletia controversa* or *Tilletia indica*), septoria disease (*Septoria nodorum*) and loose smut;
- wheat, as regards controlling the following diseases of the aerial parts of the  
 20 plant: cereal eyespot (*Tapesia yallundae*, *Tapesia acuiformis*), take-all (*Gaeumannomyces graminis*), foot blight (*F. culmorum*, *F. graminearum*), black speck (*Rhizoctonia cerealis*), powdery mildew (*Erysiphe graminis forma specie tritici*), rusts (*Puccinia striiformis* and *Puccinia recondita*) and septoria diseases (*Septoria tritici* and *Septoria nodorum*);
- 25 - wheat and barley, as regards controlling bacterial and viral diseases, for example barley yellow mosaic;
- barley, as regards controlling the following seed diseases: net blotch (*Pyrenophora graminea*, *Pyrenophora teres* and *Cochliobolus sativus*), loose smut (*Ustilago nuda*) and fusaria (*Microdochium nivale* and *Fusarium roseum*);
- 30 - barley, as regards controlling the following diseases of the aerial parts of the plant: cereal eyespot (*Tapesia yallundae*), net blotch (*Pyrenophora teres* and *Cochliobolus sativus*), powdery mildew (*Erysiphe graminis forma specie hordei*), dwarf leaf rust (*Puccinia hordei*) and leaf blotch (*Rhynchosporium secalis*);
- potato, as regards controlling tuber diseases (in particular *Helminthosporium solani*, *Phoma tuberosa*, *Rhizoctonia solani*, *Fusarium solani*), mildew (*Phytophthora infestans*) and certain viruses (virus Y);
- 35

- potato, as regards controlling the following foliage diseases: early blight (*Alternaria solani*), mildew (*Phytophthora infestans*);
- cotton, as regards controlling the following diseases of young plants grown from seeds: damping-off and collar rot (*Rhizoctonia solani*, *Fusarium oxysporum*) and black root rot (*Thielaviopsis basicola*);
- protein yielding crops, for example peas, as regards controlling the following seed diseases: anthracnose (*Ascochyta pisi*, *Mycosphaerella pinodes*), fusaria (*Fusarium oxysporum*), grey mould (*Botrytis cinerea*) and mildew (*Peronospora pisi*);
- oil-bearing crops, for example rape, as regards controlling the following seed diseases: *Phoma lingam*, *Alternaria brassicae* and *Sclerotinia sclerotiorum*;
- corn, as regards controlling seed diseases: (*Rhizopus* sp., *Penicillium* sp., *Trichoderma* sp., *Aspergillus* sp., and *Gibberella fujikuroi*);
- flax, as regards controlling the seed disease: *Alternaria linicola*;
- forest trees, as regards controlling damping-off (*Fusarium oxysporum*, *Rhizoctonia solani*);
- rice, as regards controlling the following diseases of the aerial parts: blast disease (*Magnaporthe grisea*), bordered sheath spot (*Rhizoctonia solani*);
- leguminous crops, as regards controlling the following diseases of seeds or of young plants grown from seeds: damping-off and collar rot (*Fusarium oxysporum*, *Fusarium roseum*, *Rhizoctonia solani*, *Pythium* sp.);
- leguminous crops, as regards controlling the following diseases of the aerial parts: grey mould (*Botrytis* sp.), powdery mildews (in particular *Erysiphe cichoracearum*, *Sphaerotheca fuliginea* and *Leveillula taurica*), fusaria (*Fusarium oxysporum*, *Fusarium roseum*), leaf spot (*Cladosporium* sp.), alternaria leaf spot (*Alternaria* sp.), anthracnose (*Colletotrichum* sp.), septoria leaf spot (*Septoria* sp.), black speck (*Rhizoctonia solani*), mildews (for example *Bremia lactucae*, *Peronospora* sp., *Pseudoperonospora* sp., *Phytophthora* sp.);
- fruit trees, as regards diseases of the aerial parts: monilia disease (*Monilia fructigenae*, *M. laxa*), scab (*Venturia inaequalis*), powdery mildew (*Podosphaera leucotricha*);
- vine, as regards diseases of the foliage: in particular grey mould (*Botrytis cinerea*), powdery mildew (*Uncinula necator*), black rot (*Guignardia biwelli*) and mildew (*Plasmopara viticola*);

- beetroot, as regards the following diseases of the aerial parts: cercospora blight (*Cercospora beticola*), powdery mildew (*Erysiphe beticola*), leaf spot (*Ramularia beticola*).

5 Preferably, the plant that can be protected by the method according to the present invention is potato.

10 The fungicide composition according to the present invention may also be used against fungal diseases liable to grow on or inside timber. The term "timber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists in contacting one or more compounds of the present invention, or a composition according to the invention; this includes for example direct application, spraying, dipping, injection or any other suitable means.

15

The dose of active material usually applied in the treatment according to the present invention is generally and advantageously between 10 and 1500 g/ha, preferably between 50 and 1000 g/ha for applications in foliar treatment. The dose of active substance applied is generally and advantageously between 2 and 200 g per 20 100 kg of seed, preferably between 3 and 150 g per 100 kg of seed in the case of seed treatment. It is clearly understood that the doses indicated above are given as illustrative examples of the invention. A person skilled in the art will know how to adapt the application doses according to the nature of the crop to be treated.


25 The fungicidal composition according to the present invention may also be used in the treatment of genetically modified organisms with the compounds according to the invention or the agrochemical compositions according to the invention. Genetically modified plants are plants into whose genome a heterologous gene encoding a protein of interest has been stably integrated. The expression 30 "heterologous gene encoding a protein of interest" essentially means genes which give the transformed plant new agronomic properties, or genes for improving the agronomic quality of the transformed plant.

35 The compositions according to the present invention may also be used for the preparation of composition useful to curatively or preventively treat human and animal fungal diseases such as, for example, mycoses, dermatoses, trichophyton

diseases and candidiases or diseases caused by *Aspergillus spp.*, for example *Aspergillus fumigatus*.

CLAIMS

- 5 1. A fungicidal composition comprising :  
 a) at least a pyridylmethylbenzamide derivative of general formula (I)



in which :

- 15 - R¹ may be a hydrogen atom, an optionally substituted alkyl group or an optionally substituted acyl group;  
 - R² may be a hydrogen atom or an optionally substituted alkyl group;  
 - R³ and R⁴ may be chosen independently from each other as being a halogen atom, a hydroxyl group, a cyano group, a nitro group, -SF₅, a trialkylsilyl group, an optionally substituted amino group, an acyl group, or a group E, OE or SE, in which  
 20 E may be an alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl or a heterocyclyl group each of which may optionally be substituted;  
 - p represents 0, 1, 2, 3 or 4;  
 - q represents 0, 1, 2, 3 or 4;

and its agriculturally acceptable optical and/or geometric isomers, tautomers and  
 25 addition salts with an acid or a base;  
 and

b) a chloronitrile derivative which is chlorothalonil;  
 in a compound (I) / chlorothalonil weight ratio of from 0,01 to 1.

- 30 2. A composition according to claim 1, characterised in that R¹ and R² are chosen independently from each other as being a hydrogen atom or an optionally substituted alkyl group.

3. A composition according to claim 2, characterised in that the optionally  
 35 substituted alkyl group is a methyl group or an ethyl group.

4. A composition according to claim 3, characterised in that  $R^1$  and  $R^2$  are both hydrogen atoms.
5. A composition according to any one of the claims 1 to 4, characterised in that  $R^3$  and  $R^4$  are chosen independently from each other as being a halogen atom, a hydroxyl group, a nitro group, an optionally substituted amino group, an acyl group, or a group E, OE or SE, in which E may be a alkyl, a cycloalkyl, a phenyl or a heterocyclyl group, each of which may optionally be substituted.
6. A composition according to claim 5, characterised in that  $R^3$  and  $R^4$  are chosen independently from each other as being a halogen atom, a nitro group or a halogenoalkyl group.
7. A composition according to claim 6, characterised in that the halogen atom is a chlorine atom and the halogenoalkyl group is a trifluoromethyl group.
8. A composition according to any one of the claims 1 to 7, characterised in that p and q are chosen independently from each other as being 1 or 2.
9. A composition according to claim 8, characterised in that p is 2.
10. A composition according to claim 8 or 9, characterised in that q is 2.
11. A composition according to any one of the claims 1 to 10, characterised in that the compound of general formula (I) is chosen as being
- a compound (Ia) which is 2,6-dichloro-N-{{3-chloro-5-(trifluoromethyl)-2-pyridinyl)methyl}benzamide; or
  - a compound (Ib) which is N-{{3-chloro-5-(trifluoromethyl)-2-pyridinyl)methyl}-2-fluoro-6-nitrobenzamide; or
  - a compound (Ic) which is N-{{3-chloro-5-(trifluoromethyl)-2-pyridinyl)methyl}-2-methyl-6-nitrobenzamide.
12. A composition according to any one of the claims 1 to 11, characterised in that the compound (I) / chlorothalonil weight ratio is from 0,01 to 1.

13. A composition according to claim 12, characterised in that the compound (I) / chlorothalonil weight ratio is from 0,025 to 0,5.
14. A composition according to claim 13, characterised in that the compound (I) / chlorothalonil weight ratio is from 0,05 to 0,1.
15. A composition according to any one of the claims 1 to 14, characterised in that it comprises, in addition to the active material, an agriculturally acceptable support, carrier or filler.
16. A composition according to claim 15 further comprising a surfactant.
17. A composition according to either of claims 15 and 16, comprising from 0.05% to 99% by weight of active material.
18. Method for preventively or curatively controlling phytopathogenic fungi of crops, characterised in that an effective and non-phytotoxic amount of a composition according to any one of the claims 1 to 17 is applied to the seed, the plant and/or to the fruit of the plant or to the soil in which the plant is growing or in which it is desired to grow.
19. Method according to claim 18, characterised in that the plant is potato.

### **ABSTRACT**

5 A fungicidal composition comprising at least a pyridylmethylbenzamide derivative  
of general formula (I) and a chloronitrile derivative which is chlorothalonil; in a  
compound (I) / chlorothalonil weight ratio of from 0,01 to 1.

10 Method for preventively or curatively combating the phytopathogenic fungi of crops  
by using this composition.